

excess cholesterol from the blood and takes it to the liver for disposal. A high HDL level is related to a lower risk of heart and blood vessel disease, ie the higher your HDL level the better.

Low-Density Lipoprotein (LDL) or “Bad cholesterol” is a lipoprotein (a combination of fat and protein) found in the blood. It is called “bad “ cholesterol because it picks up cholesterol from the blood and takes it to the cells. A high LDL level is related to a higher risk of heart and blood vessel disease. Reducing the LDL level is a major treatment target for cholesterol-lowering medications.

HEPATITIS B (Hep B)

Hepatitis B is a disease caused by the hepatitis B virus (HBV). Infection with this virus can cause scarring of the liver, liver failure, liver cancer, and even death. Hepatitis B is spread by infected blood and other bodily fluids such as semen, vaginal secretions, and open sores.

HIV

The human immunodeficiency virus (HIV) is spread by sexual contact with infected blood, and from mother to child during pregnancy, childbirth, or breastfeeding. HIV interferes with your body’s ability to fight the organisms that cause disease by weakening your immune system to the point that you have AIDS (acquired immune deficiency syndrome)

STOOL TESTS

These may be used to diagnose the presence or absence of several medical conditions.

1. FAECAL IMMUNOCHEMICAL TEST (FIT): This detects blood in the stool that is not visible (occult). Common causes of bleeding in the digestive tract include peptic ulcer disease, cancer of the stomach, colon, or rectum. It is not affected by intake of red meat, iron, or aspirin. Other advantages of FIT include its non-invasive nature and low cost compared to other screening methods. It is recommended annually and is commonly the test of choice in programmatic screening.

2. Microscopic examination of stools may reveal the presence of larvae or eggs of worms in the intestines. This can aid in the diagnosis of intestinal worms including Ascariasis, Hookworm, Strongyloidiasis, and Whipworm.

CERVICAL CANCER SCREENING IN WOMEN

1. Pap Smear: A Pap smear is used to test for cervical cancer in women. It involves collecting cells from your cervix – the lower, narrow end of your uterus that is at the top of your vagina. Detecting these abnormal cells early with a Pap smear is the first step in halting your possible development of cervical cancer.

2. HPV screening: The human papillomavirus (HPV) test detects the presence of human papillomavirus, the virus that causes during the pap test can be used to test for HPV at the laboratory.

PROSTATE CANCER SCREENING IN MEN

Two tests are commonly used to screen for prostate cancer :

1. Digital rectal exam (DRE): A doctor inserts a gloved, lubricated finger into your rectum to estimate the size of your prostate and feel for lumps or other abnormalities.

2. Prostate-specific antigen (PSA) test: This test measures the level of PSA in your blood. PSA is a substance made by our prostate. As a rule the higher the PSA level in your blood the more likely a prostate problem is present. PSA levels also can be affected by certain medical procedures, medications, enlarged prostate, and prostate infection. Because many factors can affect PSA levels your doctor is the best person to interpret your PSA test results.

References:

www.mayoclinic.org/test
www.WebMD.com
www.nlm.nih.gov/medlineplus
www.cdc.gov

PATHOLOGISTS LANCET KENYA

Main Laboratory / Headquarters
5th Ave. Office Suites- Upper Hill
5th Ngong Avenue , Ngong Road

Phone: 0703061000

Email: info@cerbalancetafrica.co.ke

Website: www.cerbalancetafrica.ke

Facebook: Pathologists Lancet Kenya

Twitter: @lancetkenya_plk

Instagram: @pathologists_lancet_kenya

Linkedin: Pathologists Lancet Kenya

HEALTH SCREENING FOR MEN AND WOMEN



A wellness test is a medical investigation, such as a yearly physical check-up, which is aimed at preventing health problems in a healthy person. It is in contrast to the type of medical tests one gets when there are specific symptoms that need to be diagnosed. In wellness tests, the doctor may draw blood or take urine and stool samples to check the level of certain substances in your body.

Some of the important tests a Doctor may request for an adult aged below 40 years are explained below. Your doctor will usually advise you on how frequently you will need to take these tests as this will depend on your gender and your present state of health and your risk factors.

FULL BLOOD COUNT (FBC) OR COMPLETE BLOOD COUNT (CBC)

The full blood count is the measure of the concentration of white blood cells, red blood cells, and platelets in the blood and helps detect a wide range of disorders including anemia infection, and leukemia. Red blood cells carry oxygen and low levels are seen in anemia. White blood cells fight infection and could be high if there is an infection or leukemia.

Platelets help with blood clotting. Abnormally high or low levels can be seen in anemia, bleeding disorders, iron deficiency, leukemia, etc.

KIDNEY FUNCTION TESTS

Urea measures the amount of urea that is in your blood. Healthy kidneys remove urea and other waste products from your blood. A urea test can reveal whether your urea levels are higher than normal, suggesting that your kidneys may not be working properly.

Creatinine is a chemical waste product that is produced by a muscle metabolism by eating meat. Healthy kidneys remove creatinine and other waste products from your blood. If your kidneys are not functioning properly an increased level of creatinine may accumulate in your blood.

LIVER FUNCTION TESTS

Your doctor may conduct liver function tests if you have liver disease you drink alcohol excessively you are taking medication that can harm the liver or you have the symptoms of liver or bile system disease (nausea and vomiting abdominal pain or yellow skin)

The liver filters and processes blood as it circulates through

the body. It metabolizes nutrients, detoxifies harmful substances, makes blood clotting proteins and performs many other vital functions.

The cells in the liver contain enzymes that carry out these chemical reactions. When liver cells are damaged or destroyed the enzymes in the cells leak out into the blood where they can be measured by blood tests. The two main enzymes that are elevated in liver diseases are aspartate aminotransferase (AST) and gamma-glutamyl transferase (GGT) may also be increased if there is an obstruction in the liver although there may be other reasons for these enzymes being increased as well.

Albumin is the main protein made by the liver. It performs many important functions including the transport of hormones, vitamins, and other substances throughout the body. A low albumin level indicates that the liver is not functioning properly. Albumin may also be decreased when you have an infection or inflammation.

Bilirubin is a waste product resulting from the breakdown of red blood cells. It is processed by the liver before being excreted through the stool. If your liver is damaged it cannot properly process bilirubin. This will lead to an abnormally high level of bilirubin in the blood.

FASTING BLOOD SUGAR

This measures blood sugar (glucose) after you have not eaten for at least 8 hours. It is often the first test done to check for diabetes. Glucose comes from carbohydrate foods and is the main source of energy used by the body. Insulin is a hormone produced in the pancreas that helps your body cells use glucose. Normally your blood glucose levels increase slightly after you eat. This increase can cause your pancreas to release insulin so that your blood glucose levels do not get too high. In diabetes, the blood glucose levels are higher than normal.

HAEMOGLOBIN A1c (HbA1c)

The haemoglobin A1c (HbA1c) test or glycated haemoglobin test is a blood test used to diagnose diabetes and to determine how well your diabetes is being controlled. Hemoglobin is a substance within red blood cells that carries oxygen throughout your body. When your diabetes is not controlled sugar builds up in your blood and combines your hemoglobin which becomes “glycated “. Haemoglobin A1c is a measure of the average concentration of your blood sugar over the preceding six to twelve-week period and is

used in conjunction with blood sugar monitoring to make adjustments in your diabetes medicines. If your blood sugar levels have been high over recent weeks your HbA1c test result will be higher than normal.

LIPID PROFILE

This test measures the number of lipids (cholesterol and other fats) in the blood and it is done to assess your risk of developing cardiovascular disease (heart and type blood vessel disease).

Excessive lipids in the blood cause a build-up of plaques in the arteries that can lead to narrowed or blocked arteries throughout the body (atherosclerosis) causing reduced blood flow.

Atherosclerosis can result in heart disease kidney disease and stroke.

Typically you will be required to fast for 10-12 hours (no food or drink except water) before the test to eliminate the contribution of any fat you recently ate.

Triglycerides (TG) are a type of fat in the blood. When you eat your body converts any calories it doesn't need into triglycerides which are stored in fat cells. High triglycerides are associated with several factors including being overweight, eating too many sweets or drinking too much alcohol, thy, smoking, a sedentary lifestyle, diabetes thyroid or liver disease, and genetic conditions. High levels of triglycerides are related to a higher risk of heart and blood vessel disease.

Total cholesterol (TC) cholesterol is a type of fat found in your blood. It is produced by our body and also comes from the foods you eat (animal products). Cholesterol is needed by our body to maintain the health of our cells. Total cholesterol level is the sum of all the types of cholesterol in your blood including low-density lipoproteins (LDL) high-density lipoproteins (HDL) and very-low-density lipoproteins (VLDL). The higher your total cholesterol the greater your risk for heart disease.

As cholesterol and triglycerides are insoluble in blood, they need to be transported in our bodies. The role of lipoprotein particles is to transport triglycerides and cholesterol in the blood between all the tissues of the body.

High-Density Lipoprotein (HDL) or “Good cholesterol” is a lipoprotein (a combination of fat and protein) found in the blood. It is called ‘good’ cholesterol because it removes